## **Tobacco-Related Cancer Deaths, Excluding Lung Cancer, by Sex, Maine, 2008**

	Number of Deaths	Crude Rate		Age-adjusted Rate	
		Rate	95% CI	Rate	95% CI
Maine Total	542	41.1	37.7 - 44.7	32.6	29.9 - 35.5
Female	206	30.5	26.5 - 35.0	21.9	19.0 - 25.3
Male	336	52.2	46.7 - 58.0	46.5	41.5 - 51.8

## **Tobacco-Related Cancer Deaths, Excluding Lung Cancer, by Age and Race, Maine, 2004-2008**

	Count	Crude Rate		Age-adjusted Rate	
	Count	Rate	95% CI	Rate	95% CI
Age					
00-14 years	NSD	NSD	NSD	NApp	NApp
15-19 years	NSD	NSD	NSD	NApp	NApp
20-24 years	NSD	NSD	NSD	NApp	NApp
25-29 years	NSD	NSD	NSD	NApp	NApp
30-34 years	NSD	NSD	NSD	NApp	NApp
35-39 years	11	2.5	1.3 - 4.5	NApp	NApp
40-44 years	44	8.6	6.2 - 11.5	NApp	NApp
45-49 years	89	16.1	13.0 - 19.9	NApp	NApp
50-54 years	151	28.9	24.4 - 33.8	NApp	NApp
55-59 years	257	55.1	48.5 - 62.2	NApp	NApp
60-64 years	323	90.2	80.6 - 100.6	NApp	NApp
65-69 years	313	115.9	103.5 - 129.5	NApp	NApp
70-74 years	402	176.7	159.9 - 194.9	NApp	NApp
80-84 years	417	284.9	258.2 - 313.6	NApp	NApp
85+ years	439	334.8	304.2 - 367.6	NApp	NApp
Race					
White	5,505	43.5	42.3 - 44.6	36.3	35.3 - 37.2
Black	11	8.7	4.3 - 15.5	26.9	13.0 - 47.6
Other*	33	16.2	11.1 - 22.7	30.7	19.6 - 45.1

Data Source: Surveillance, Epidemiology, and End Results SEER Program, National Cancer Institute, Underlying mortality data provided by National Center for Health Statistics. Rates are calculated using SEER\*Stat Version 7.0.5.

Cancer Deaths: Deaths with malignant cancer as the underlying cause of death.

Tobacco related Cancer excluding Lung: laryngeal, oropharyngeal, esophageal, stomach, pancreatic, kidney and renal pelvis, urinary bladder, cervical cancers, and acute myeloid leukemia. SEER Cause of Death Recode: 22020, 20080,21100 ,21010,21020,29020,29010,27010, 3502 (which include 1999-2009: ICD-10 codes C32,C10,C15-C16, C25,C53,C64-C65,C67,C92.0, C92.4-C92.5, C94.0, C94.2;1993-1998: ICD-9 codes 161,146.3-146.9,150-151,157,180,188,189.0-189.1,205.0, 207.0, 207.2). Crude rates are deaths per 100,000 population. Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population. 95% CI: 95% confidence interval of the rate.

NSD: Non-sufficient data. Data is not available due to small or zero counts.

Napp: Not applicableage-adjusted rates are not calculated for age-specific groups.

<sup>\*</sup>Other Race includes American Indian/AK Native and Asian/Pacific Islander; NSD = Non sufficient data. Data not available due to too few cases.